

Spin fractionalization of an even number of electrons in a quantum dot

Domenico Giuliano^{1,2} and Arturo Tagliacozzo^{1,3}

¹ *Istituto Nazionale di Fisica della Materia (INFM), Unità di Napoli*

² *Department of Physics, Stanford University, Stanford, California 94305*

³ *Institut für Nanotechnologie (INT), Forschungszentrum, Postfach 3640, D-76021 Karlsruhe, Germany
and Dipartimento di Scienze Fisiche Università di Napoli "Federico II",
Mostra d' Oltremare Pad. 19, I-80125 Napoli, Italy*

Kondo resonant conductance can occur in a quantum dot with an even number of electrons N in the Coulomb blockade regime, with bottom and top contacts attached to it in a pillar configuration, in a magnetic field $B_{\perp} = B^*$ along the axis. The field B^* is tuned at the crossing of levels with different total spins and the spin degeneracy of the dot energies should be lifted. The case of the degeneracy of the singlet-triplet states in a dot with $N = 2$ is analyzed in detail. Coupling to the contacts is antiferromagnetic due to a spin selection rule, so that total spin on the dot is $S = 1/2$ in the Kondo state.

Quantum dots (QD) are remarkable because of Coulomb oscillations [1]. At very low temperature, linear conductance (at vanishing bias V_{sd}) is zero, except for peaks at discrete values of the gate voltage V_g , when it is energetically favourable to add one extra electron to the dot. Therefore V_g controls the number of particles on the dot in the Coulomb Blockade (CB) regime: its charge is quantized and so is the spin. Dots with an even (odd) number of electrons have integer (half integer) total spin.

However, by lowering the temperature further below some value T_K , finite conductance can be found experimentally at zero voltage [2-4], relatively insensitive to small changes of V_g within the CB regime, as a mesoscopic realization of the physics of the Kondo effect [5-7]. The zero voltage anomaly in the conductance is only seen when population is odd [3]. While the charge remains quantized, the spin quantization of the correlated state between the dot and its contacts is changed, due to the formation of a Kondo singlet involving the odd electron.

A weak magnetic field parallel to the dot plane, $B_{\parallel}$, splits the peaks in the differential conductance into two, located at $V_{sd} = \pm g^* \mu_B B_{\parallel}$, (where g^* is the effective giromagnetic factor for electrons in this geometry and $\mu_B = e\hbar/2mc$ is the Bohr magneton), until eventually they disappear [3,4].

On the contrary, a magnetic field $B_{\perp}$, orthogonal to the dot area, is believed to be much more destructive, due to orbital changes of the electron states [8], in addition to the lifting of spin degeneracy, that prevents the formation of the Kondo correlated state.

We suggest here the possibility that, due to accidental crossing of the lowest non degenerate many body energy levels of the QD at special values of $B_{\perp} = B^*$, orthogonal to the dot connected to the leads in a pillar structure, the Kondo resonance can still be present, irrespective of the parity of the electron number N [9,10]. In particular we discuss here the case of $N = 2$ and show that under suitable conditions a correlated state sets in, which is

characterized by fractionalization of the total spin.

Indeed a QD at high $B_{\perp}$ (in the z -direction) displays a rich level structure that can be monitored attaching bottom and top contacts to it in a pillar structure via linear and non linear transport [11-14]. Assuming the QD to be effectively two-dimensional and confined by a parabolic potential, its states in isolation can be labelled by the total spin S and the z -component of the total angular momentum M and spin S_z . In increasing $B_{\perp}$, many level crossings can occur for an N particle QD between states of increasing M , as well as states with higher total spin S .

It is well known that the ground state (GS) of an isolated dot with $N = 2$ electrons changes from singlet ($^2S : S = 0, M = 0$) to triplet ($^2T_{S_z} : S = 1, M = 1$) at a value $B_{\perp} = B^*$, due to crossing of levels [15]. Correspondingly, exact diagonalization shows that the $N = 3$ electron dot is in a doublet state ($^3D_{S_z} : S = 1/2, M = 1$ or 2 depending on the field value) (see fig.1) [14]. As shown quantitatively in the following, the dot parameters can be chosen in such a way that $B_{\perp}$ is high enough to lift spin degeneracy. We take V_g tuned in the CB region between $N = 2$ and $N = 3$ electrons on the dot, and limit our discussion to four states : 1) dot with $N = 1$ in 1D , 2-3) dot with $N = 2$ in one of the two degenerate states 2S_0 , and 2T_1 , 4) dot with $N = 3$ in $^3D_{1/2}$. Taking only these states allows the mapping of the problem onto the non degenerate Anderson model with the following levels: 1) empty level of zero energy, 2-3) singly occupied level of energy ϵ_d (doubly degenerate), 4) doubly occupied level of energy $2\epsilon_d + U$. Here U is very large (see below) and, if levels 2-3 and 4 are symmetric with respect to the chemical potential of the contacts μ , what can be controlled by V_g , average occupancy of levels 2-3 is $n_d = 1$. This corresponds to $N = 2$ electrons on the dot.

The key observation is the following spin selection rule for the isolated dot [16] : because the $N = 3$ GS for the dot only has $S_z = 1/2$, hybridization of the dot with the contacts can only occur by virtual occupancy of the

dot with an extra spin up electron when the dot is in the $N = 2$ singlet state 2S or with an extra spin down electron when the dot is in the $N = 2$ triplet state with $S_z = 1$ (2T_1 state). Although spin degeneracy of the dot is assumed to be lifted, the physics is the same as that of the single channel Kondo model with effective spin $S_z^{eff} = 1/2$, in the absence of magnetic field. Here is $S_z^{eff} = S_z - 1/2$ and coupling with the delocalized electrons is antiferromagnetic due to the spin selection rule mentioned above. Provided hybridization to the contacts Δ is large enough and temperature is below the Kondo temperature T_K , the system flows to the strongly correlated fixed point. Formation of the Kondo singlet corresponds in this case to an average spin on the dot frozen at $S_z = \frac{1}{2}$, that is to fractionalization of the total spin of the dot, together with a strong zero bias anomaly in the conductance.

In this letter we discuss the properties of the GS, borrowing from the known results of the non degenerate Anderson model [17]. In particular, we discuss the spin fractionalization, the practical realizability of the system and the robustness of the GS versus small changes of the magnetic field $B_\perp$ with respect to the value B^* that makes the states $|S\rangle$ and $|T\rangle$ degenerate in energy.




FIG. 1. Calculated energies of a circular quantum dot vs magnetic field $B_\perp$ (expressed in cyclotron energies $\hbar\omega_c$) for $N = 2$ (a),(c) and $N = 3$ (b),(d). Here the level spacing of the confining potential is $\hbar\omega_d = 4\text{meV}$ and the strength of the Coulomb interaction is $E_c = 3\text{meV}$ (notations can be found in the text). (c),(d) magnify the marked area of (a),(b), respectively [14]

We denote the two dot degenerate energies by 2E_i ($i = 1$: triplet state, $i = 2$: singlet state) and take the chemical potential μ of the left and right bulk con-

tacts within the CB valley of the conductance at $N = 2$ ($\mu_N < \mu < \mu_{N+1} \equiv {}^{N+1}E - {}^N E_i$). Electrons in the contacts are non interacting and have an energy dispersion $\epsilon_{k\sigma}$ involving just one single Landau level and linearized around μ with Fermi velocity $v_{F\sigma}$. The wavevector k is in the z - direction and the two contacts are assumed to be equal ($H_{L/R} = \sum_{k\sigma} \epsilon_{k,\sigma} b_{L/R,k\sigma}^\dagger b_{L/R,k\sigma}$) and three dimensional enough, in order to have both spin polarizations at μ .

The dot Hamiltonian H_d , restricted to the Fock space with $N = 1, 2, 3$ electrons can be represented in terms of the occupation numbers n_i for the degenerate levels 2E_i :

$$H_d = \mu_2 \sum_i n_i + U n_1 n_2 \quad (1)$$

where $U = \mu_3 - 2\mu_2$ is the charging energy for adding the third electron to the dot. Here is $n_i = c_i^\dagger c_i$, the vacuum of the c operators is the dot state with one electron and the $c^\dagger$ create the two degenerate states by acting on it.

The model Hamiltonian is $H = H_L + H_R + H_d + H_t$, where H_t is the hybridization term acting only at $z = 0$. According to the spin selection rule stated above, it takes the form:

$$H_t = \frac{1}{\sqrt{2}} \sum_k \left\{ \Gamma^* (b_{L,k\uparrow}^\dagger + b_{R,k\uparrow}^\dagger) c_1 + \Gamma c_2^\dagger (b_{L,k\downarrow} + b_{R,k\downarrow}) \right\} + h.c.$$

The tunneling amplitudes $\Gamma_{L/R,k\sigma i}$, which depend on the actual features of the device, have been taken all equal for simplicity.

We start integrating out the contacts fields ($\hbar = 1$ in the following):

$$\mathcal{Z}(\mu) \propto \int \Pi_i \left(D c_i D c_i^\dagger \right) e^{-\mathcal{A}_d} e^{-\beta |\Gamma|^2 \sum_{\omega_m, i} c_i^\dagger K(i\omega_m) c_i} \quad (2)$$

with

$$\mathcal{A}_d = \int_0^\beta d\tau \left\{ \sum_i \left[c_i^\dagger \frac{\partial}{\partial \tau} c_i + (\mu_2 - \mu) c_i^\dagger c_i \right] + U c_1^\dagger c_1 c_2^\dagger c_2 \right\}$$

and $K(\omega_m) = \sum_k (i\omega_m + vk)^{-1}$ is taken independent of σ for simplicity. In the limit of large U we have:

$$\exp \left[\int_0^\beta d\tau [U n_1 n_2 - \tilde{\mu} (n_1 + n_2)] \right]$$

$$= e^{\frac{\beta}{4U} \tilde{\mu}^2} \delta(n_1 + n_2 - 2\tilde{\mu}/U) \cdot e^{-\frac{U}{4} \int_0^\beta d\tau (n_1 - n_2)^2},$$

where the delta function implements the constraint of single site occupancy in the symmetric case, $\tilde{\mu} = \mu - \mu_2 = U/2$.

We introduce $X(\tau)$ as an auxiliary boson field coupled to the differential occupancy of the two degenerate states by means of an Hubbard- Stratonovitch decoupling:

$$e^{\frac{U}{4} \int_0^\beta d\tau (n_1 - n_2)^2} =$$

$$\int DX e^{-\frac{1}{4U} \int_0^\beta d\tau (X^2(\tau) + 2U(n_1 - n_2)X(\tau))}.$$

Then, the partition function of eq.(2) reads:

$$\begin{aligned} \mathcal{Z}(\mu) &\propto \int DX e^{-\frac{1}{4U} \int_0^\beta d\tau X^2(\tau)} \\ &\times \int \Pi_i \left(Dc_i Dc_i^\dagger e^{-\int_0^\beta d\tau d\tau' c_i^\dagger(\tau) G_{(0)}^{-1}(\tau - \tau') c_i(\tau')} \right) \\ &\times \delta(n_1 + n_2 - 1) \sum_{j=1,2} e^{(-1)^j \int_0^\beta d\tau [n_j - \frac{1}{2}] X(\tau)}, \quad (3) \end{aligned}$$

where is $G_{(0)}^{-1}(i\omega_m) = i\omega_m + |\Gamma|^2 K(i\omega_m)$. We will take the saddle point solution of eq.3. At saddle point (see below) the constraint is satisfied in the average over the time β because $\langle n_i - \frac{1}{2} \rangle \approx -(-1)^i \overline{X_{s.p.}(\tau)} / 2U = 0$ and the integrals for $i = 1, 2$ decouple, giving a contribution to the effective action for the field X that, for $i = 1$, is given by $\int_0^1 dg \int_0^\beta d\tau [G_{(gX),1}(\tau, \tau^+) - \frac{1}{2}] X(\tau)$ with $G_{(gX),1}^{-1} = G_{(0)}^{-1} - gX$ ($X \rightarrow -X$ for $1 \rightarrow 2$) [18]. This representation allows to recognize the doubly degenerate GS of an effective spin $S_z^{eff} = 1/2$ with $S_z^{eff} = (S_z - 1/2)$, driven by the field $X(\tau)$ and produced by virtual tunneling of electrons on and off the dot at energy μ .

We now follow ref. [18] quite closely, to map eq.3 on a 1-dimensional Coulomb gas (CG). We define $\xi(\tau) = X(\tau)/\Delta$, where $\Delta = 2\pi N(0)\Gamma^2$ and $N(0) = L/\pi v_F$ (L is the size of each single contact) is the density of states at the Fermi energy. At low temperatures, saddle point solutions of the resulting single particle effective action are sequences of instantons $\xi_\pm(\tau, l) = \pm \xi_0 \tanh((\tau - t_l)/\tau_0)$ (t_l are the centers, $l = 1, 2, \dots$) corresponding to jumps between the two minima of the effective potential, $V[\xi] = \Delta^2/U \xi^2 - 2\Delta/\pi [\xi \text{tg}^{-1}(\xi) - \frac{1}{2} \ln(1 + \xi^2)]$ [18], located at $\xi_0 = \pm U/2\Delta$, that interact via a logarithmic potential $\alpha^2 \ln |(t_i - t_j)/\tau_0|$.

The bare strength of the logarithmic interaction is $\alpha_b^2 = 2(1 - 4\Delta/U\pi^2)$. The bare fugacity of the CG is [19] $Y_b = \tau_0 \exp(-\bar{\mathcal{A}})$ ($\bar{\mathcal{A}} \sim \tau_0 U$ is the action of one single blip). The scaling of the fugacity and the renormalization of the coupling constant induced by processes of fusion of charges lead to the Renormalization Group (RG) equations [20]:

$$\frac{d\bar{Y}}{d \ln \tau_0} = (1 - \frac{\alpha^2}{2}) \bar{Y}; \quad \frac{d\alpha^2}{d \ln \tau_0} = -2Y^2 \alpha^2 \quad (4)$$

The flow is towards $Y \rightarrow \infty$ and $\alpha^2 \rightarrow 0$ and the scaling invariant energy is $T_K = \tau_0^{-1} e^{1/(1 - \frac{\alpha^2}{2})} \sim (U\Delta)^{\frac{1}{2}} e^{-\pi^2 U/(4\Delta)}$. Condensation of instantons in the doubly degenerate GS leads to the Kondo singlet $\langle S_z^{eff} \rangle = 0$ or $\langle S \rangle = \frac{1}{2}$ on the dot.

An heuristic argument for the fractionalization of the total spin S of the dot reads as follows.

The dynamics of the field $\xi(\tau)$ of action $\bar{\mathcal{A}}$ can be mimicked by a two level system hamiltonian H_{2l} with hopping energy $\lambda \sim \frac{1}{2} m_{eff} \left(\frac{d\xi}{d\tau} \right)^2 \sim \frac{1}{2} \frac{\Delta^2}{U} \tau_0^2 (\xi_0/\tau_0)^2 \sim U/2$. The role of the interaction is to project out higher energy components from the dot states. Denoting by $|\pm \rangle$ the two eigenstates of H_{2l} , the dynamics of the dot is between states $\Pi^g |S \rangle / \mathcal{N}_S^{\frac{1}{2}}(g)$ and $\Pi^g |T \rangle / \mathcal{N}_T^{\frac{1}{2}}(g)$, where $|S \rangle$ and $|T \rangle$ are the states localized in the two potential wells and $\mathcal{N}_{S/T}(g)$ are their normalizations. $\Pi^g = 1 - g\Pi^-$ is the Gutzwiller operator partly eliminating the component on the high energy eigenstate $|- \rangle$ ($g \in (0, 1)$) [21].

Now we state the connection with the CG picture given above. The frequency $\lambda/2\pi$ is obviously related to the average number of flips: $\beta\lambda/2\pi = \langle N \rangle = Y \tanh Y$ and probability of having the system in states $|\pm \rangle$ at temperature β^{-1} are

$$P_- = \frac{e^{-4\pi Y \tanh Y}}{1 + e^{-4\pi Y \tanh Y}}; \quad P_+ = \frac{1}{1 + e^{-4\pi Y \tanh Y}}. \quad (5)$$

Equating these probabilities to the ones given by $P_\pm = |\langle \pm | \Pi^g | S/T \rangle|^2 / \mathcal{N}_{S/T}(g)$ yields the correspondence $1 - g \rightarrow e^{-2\pi Y \tanh Y}$.

The z - component of the total spin of the projected singlet and triplet state are given by:

$$\langle S_z \rangle_{S,g} = \frac{g^2}{2} \frac{1}{1 + (1 - g)^2}; \quad \langle S_z \rangle_{T,g} = 2 \frac{(1 - g/2)^2}{1 + (1 - g)^2}.$$

Because Y scales to infinity, $g \rightarrow 1$: the higher energy state completely decouples and $\langle S_z \rangle_{S/T,g} \rightarrow \frac{1}{2}$.

We now give some estimates for a possible practical realization.

If the level separation of the parabolic confining potential of the dot is $\hbar\omega_d = 4meV$, the effective frequency in presence of $B_\perp$ is $\omega_0 = \sqrt{\omega_d^2 + \omega_c^2/4}$ ($\omega_c = eB_\perp/m^*c$ is the cyclotron frequency with $m^* = 0.067m_e$ for $GaAs$). The Coulomb interaction between electrons in the plane is $E_c \times l_0/|\vec{r} - \vec{r}'|$ (where $E_c = 3meV$ and $l_0^2 = \hbar/m^*\omega_0$). This choice of the parameters gives a singlet-triplet (S-T) transition for the dot with $N = 2$ at $B_\perp = B^* \approx 4.5$ Tesla, close to one of the Delft experiments [11]. Fig.1 shows the total energy of the dot *vs* magnetic field for $N = 2$ and $N = 3$. Zeeman spin splitting of the energies is *not* included in the picture for clarity, although it is sizeable at these fields: $g^* \mu_B B = 0.11meV$. Here we assume the bulk giromagnetic factor for $GaAs$ $g^* = -0.44$, but it could be lower [22,23]. In any case, at temperatures below $300mK$ the spin degeneracy is lifted. According to fig.1, the energy difference between the GS's when three and two electrons are on the dot is $U = {}^3E - {}^2E = 12.5meV$ at the S-T crossing, which is about 1/3 higher than the value

$E_c + \hbar\omega_0 = 3\text{meV} + 5.55\text{meV}$ predicted by a constant interaction model. The $N = 3$ particle state is a doublet ($S = 1/2$) 3D with $M = 1$. Higher magnetic fields produce its crossing with the doublet with $M = 2$, while, increasing $B_\perp$ further, the quadruplet state ($Q : S = 3/2$) becomes lower in energy.

The order of magnitude of the hybridization of the dot with the contacts Δ is fixed by the requirement that neighbouring CB peaks in the conductance do not overlap [3]. Their separation is $\sim \hbar\omega_0$. For values of $\Delta \sim (0.1 - 0.2)\hbar\omega_0$, T_K (given below) is of tenths of mK . We have neglected the excited states of the two particle dot in the description. This is allowed because T_K is much less than the spin splitting of energy levels. The next available state in the spectrum of the isolated dot with $N = 2$ is the triplet state with $S_z = 0$ (2T_0). Because it is non degenerate, no Kondo-like coupling is possible. This state can only hybridize with conduction electrons via cotunneling processes which, in the elastic case, give rise to currents that are linear in the biasing voltage V_{sd} and of the order of $\sim (\Delta^2/U) V_{sd}$.

We conclude that the problem is equivalent to a non degenerate Anderson model in absence of magnetic field. At $B_\perp = B^*$, the $N = 2$ particle dot can be in the 2S_0 or 2T_1 state, but addition of one more electron requires an extra energy U . Furthermore is $U \gg \Delta$. The results are expected to be quite robust with respect to small changes of the magnetic field away from B^* , especially for $B_\perp < B^*$ because the reference field $H_0 = (U\Delta)^{1/2}/\mu_B$ is of the order of *Tesla*. For $B_\perp > B^*$, instead, the state 2T_1 will be eventually favoured in energy.

A small a.c. magnetic field $|B_\parallel| \ll |\delta B| \sim T_K/\mu_B$ (with $\delta B = B_\perp - B^* < 0$) could induce transitions between the two degenerate Kondo GS's, with Rabi frequency $\hbar\omega_R = M_d|\delta B|$. The quantity $M_d/2 \approx \frac{1}{2}g^*\mu_B[1 + \delta B/(\pi T_K)]$ is the expected magnetic moment of the Dot with $N = 2$, in the Kondo state. The measurement, if feasible, should be performed just off $B_\perp = B^*$ for $\delta B < 0$, not to match the energy separation of the two levels 2T_1 and 2T_0 .

In conclusion, we have discussed the possibility that the total spin S of a dot in CB with $N=2$ is strongly coupled to the spins of the electrons in the leads. This kind of mesoscopic Kondo effect would give rise to fractionalization of S in the dot, i.e. to a sort of 'spinon box'.

The authors acknowledge useful discussions with B.Altshuler, G.Falci, L.Glazman, F.Hekking, B.Jouault, B.Kramer, G.Morandi, H.Schöller, J.Schmid and J.Weis. Work supported by INFM (Pra97-QTMD) and by EEC with TMR project, contract FMRX-CT98-0180.

- [1] L.P. Kouwenhoven *et al.*, in "Mesoscopic electron transport", NATO ASI Series **E 345**,105; L.Sohn, L.P.Kouwenhoven and G.Schön eds., Kluwer, Dordrecht, Netherlands (1997)
- [2] D.Goldhaber-Gordon, H.Shtrikman, D.Mahalu, D.Abusch-Magder, U.Meirav and M.A.Kastner, *Nature* **391**, 156 (1998)
- [3] S.M.Cronenwett, T.H.Oosterkamp and L.P.Kouwenhoven *Science* **281**,540 (1998)
- [4] J.Schmid, J.Weis, K.Eberl and K.v.Klitzing, *Physica B* **256-258**,182 (1998)
- [5] L.I.Glazman and M.E.Raikh, *Pis'ma Zh.Eksp.Teor.Fiz.* **47**,378 (1988) [*JETP Lett.* **47**,452 (1988)];
- [6] T.K. Ng and P.A.Lee, *Phys.Rev.Lett.* **61**,1768 (1988)
- [7] Y.Meir, N.S.Wingreen and P.A.Lee, *Phys.Rev.Lett.* **70**, 2601 (1993)
- [8] W.Xue and P.A.Lee *Phys.Rev.* **B38**,3913 (1988)
- [9] The same idea but in a different geometry (with a magnetic field parallel to the dot) and a quite different model is in: M.Pustilnik, Y.Avishai and K.Kikoin *cond-mat/9908004 v2*
- [10] Most likely a split gate QD with $B_\perp$ orthogonal to the plane would have fully spin polarized Hall states in the contacts, at least in the integer QH case. (I.V.Kukushkin, K.v.Klitzing and K.Eberl *Phys.Rev.* **B55**, 10607 (1997)). It will be apparent in the following that electrons in the contacts with both spins are required for spin fractionalization. However, Kondo transmission could still take place in this case (D.Giuliano and A.Tagliacozzo, *Proc. of the 6th Int. Conf. "Path integrals from meV to TeV"*, Florence, August 25-29,1998; pg.459, R.Casalbuoni, R.Giachetti, V.Tognetti, R.Vaia, P.Verrucchi eds., World Scientific, Singapore (1999)).
- [11] L.P.Kouwenhoven *et al.* *Science* **278**,1788 (1997)
- [12] S.Tarucha *et al.*, *Phys. Rev. Lett.* **77**,3613 (1996).
- [13] B.Jouault *et al.* *cond-mat/9810094*, to appear in *Europhys. J.*
- [14] B.Jouault, G.Santoro, A.Tagliacozzo accepted on *Phys. Rev. B*: eigenenergies of up to 7 electrons in a QD are obtained using a Lanczos code with a basis of 28 single particle orbitals.
- [15] M.Wagner, U.Merkt, A.V.Chaplik, *Phys.Rev.* **B45**, 1951 (1992)
- [16] D.Weinmann, W.Häusler, W.Pfaff, B.Kramer and U.Weiss, *Europhys. Lett.* **26**,467 (1994)
- [17] A.M.Tsvelick and P.B.Wiegmann, *Advances in Physics* **32**, 453 (1983)
- [18] D.R.Hamann, *Phys.Rev.* **B2**,1373 (1970)
- [19] P.W.Anderson, G.Yuval and D.R.Haman *Phys.Rev.* **B11**, 4464 (1970)
- [20] B.Nienhuis, "Coulomb Gas Formulation of Two-dimensional Phase Transitions", C. Domb and J.Lebowitz Eds., Academic Press (1987)
- [21] F. Gebhard, D. Vollhardt, *Phys. Rev. Lett.* **59**, 1472 (1987)
- [22] M.Dobers, K.v.Klitzing and G.Weinmann, *Phys.Rev.* **B38**, 5453 (1988)
- [23] note that with $\hbar\omega_d = 8\text{meV}$ and $E_c = 6\text{meV}$ used in ref [13] the S-T crossing takes place at 11*Tesla* with a spin splitting of $g^*\mu_B B = 0.27\text{meV}$.